



# Specimen Label

## Harvester®

### Landscape and Aquatic Herbicide

For use as a general herbicide to control weeds in commercial greenhouses and nurseries; ornamental seed crops\*; landscape, industrial, recreational, commercial, residential, and public areas; turf renovation; dormant established turfgrass; and aquatic areas.

#### Active Ingredient

Diquat dibromide: [6,7-dihydrodipyrido(1,2-a:2',1'-c)pyrazinediium dibromide].....37.3%

Other Ingredients.....62.7%

**Total** .....100.0%

Contains 2 pounds diquat cation per one (1) U.S. gallon (3.73 pounds diquat dibromide per gallon).

### Keep Out of Reach of Children

## CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, **read *Warranty Disclaimer* and *Misuse statements* inside label booklet.** If terms are unacceptable, return at once unopened.

FIRST AID	
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call <b>INFOTRAC at 1-800-535-5053.</b>	

\*Except in the state of California

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if inhaled. Harmful if swallowed. Causes moderate eye irritation. Avoid breathing spray mist and contact with eyes or clothing.

**Note to Physicians:** To be effective, treatment for diquat poisoning must begin **IMMEDIATELY**. Treatment consists of binding diquat in the gut with suspensions of activated charcoal or bentonite clay, administration of cathartics to enhance elimination, and removal of diquat from the blood by charcoal hemoperfusion or continuous hemodialysis.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils.

#### Mixers, Loaders, Applicators and Other Handlers Must Wear:

- Coveralls over long-sleeved shirt and long pants;
- Chemical-resistant gloves;
- Chemical-resistant footwear plus socks;
- Protective eyewear;
- Chemical-resistant headgear for overhead exposure;
- Chemical-resistant apron when cleaning equipment, mixing, or loading; and
- Face shield when mixing or loading.

*Exception:* After this product has been diluted to 0.50% or less in water (i.e., the labeled rate for some spot applications), applicators for *AQUATIC SURFACE APPLICATIONS* must, at a minimum, wear (Note: Mixers and loaders for this application method must still wear the PPE as described in the above section.):

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Waterproof gloves; and
- Protective eyewear.

*Exception:* At a minimum, applicators for *AQUATIC SUBSURFACE APPLICATIONS* must wear (Note: Mixers and loaders for this application method must still wear the PPE as described in the above section.):

- Short-sleeved shirt and short pants;
- Waterproof gloves; and
- Chemical-resistant footwear plus socks.

### USER SAFETY REQUIREMENT

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROLS

Mixers and loaders supporting aerial applications are required to use closed systems that provide dermal protection. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR

170.240(d)(4)]. When using the closed system, mixers and loaders' PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### **USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to aquatic invertebrates. For Terrestrial Uses do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. For Aquatic Uses do not apply directly to water except as specified on this label.

---

### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read the entire label. Use strictly in accordance with precautionary statements and directions for use, and with applicable state and federal regulations.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system. Do not use this product for reformulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the WPS.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.**

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeve shirt and long pants;
- Chemical-resistant gloves;
- Chemical-resistant footwear plus socks;
- Protective eyewear; and
- Chemical-resistant headgear for overhead exposure.

## **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Keep all unprotected persons out of operating areas or vicinity where there may be drift.**

**For terrestrial uses**, do not enter or allow entry of maintenance workers into treated areas, or allow contact with treated vegetation wet with spray, dew, or rain, without appropriate protective clothing until spray has dried.

**For aquatic uses**, do not enter treated areas while treatments are in progress.

## **PRODUCT INFORMATION**

This product is a nonvolatile herbicide for general weed control in:

- Commercial greenhouses and nurseries;
- Ornamental seed crops (flowers, bulbs, etc. - except in the state of California);
- Landscape, industrial, recreational, commercial, residential, and public areas;
- Turf renovation (all turf areas except commercial sod farms);
- Dormant established turfgrass (Bermudagrass, zoysiagrass, nonfood or feed crop); and
- Aquatic areas.

Absorption and herbicidal action is usually quite rapid with effects visible in a few days. This product controls weeds by interfering with photosynthesis that occurs within green plant tissue. Weeds should be succulent and/or actively growing for best results.

Rinse all spray equipment thoroughly with water after use. Avoid spray drift to crops, ornamentals, and other desirable plants during application, as injury may result. Application to muddy water may result in reduced control. Minimize creating muddy water during aquatic application. Use of dirty or muddy water for diluting this product may result in reduced herbicidal activity. Avoid applying under conditions of high wind, water flow, or wave action.

## **Spray Drift Management**

Avoiding spray drift at the application site is the responsibility of the applicator. The interactions of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops:

- The distance of the outermost nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor; and
- Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they must be observed.

## **Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See *Wind, Temperature and Humidity*, and *Temperature Inversions* sections of this label).

## **Controlling Droplet Size**

**Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

**Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

**Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

## **Boom Length**

For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

## **Application Height**

Applications should not be made at a height greater than 10 feet above the top of the target plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

## **Swath Adjustment**

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

## **Wind**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

## **Temperature and Humidity**

When making applications in low relative humidity conditions set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

## **Temperature Inversions**

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to

remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### **Sensitive Areas**

The pesticide should only be applied when the wind is blowing away from adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops).

### **USE IN COMMERCIAL GREENHOUSES AND NURSERIES**

For general weed control in commercial greenhouses (beneath benches), field grown and container stock, and other similar areas, this product may be applied before crop emergence either pre- or post-plant in field grown ornamental nursery plantings, or post-emergence as a directed spray. This product may also be applied before crop emergence in ornamental seed crops (except in the state of California).

Avoid contact with desirable foliage as injury may occur. Do not use on food or feed crops.

**Spot spray:** Apply 1-2 quarts of this product plus the labeled rate of a 75% or greater nonionic surfactant per 100 gallons of water, or 0.75 ounces (22 milliliters) of this product plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon of water.

**Broadcast:** Apply 1-2 pints of this product in a minimum of 15 gallons of water per acre. Add the labeled rate of a 75% or greater nonionic surfactant per 100 gallons of spray mixture. Use an adequate spray volume to insure good coverage.

### **USE IN ORNAMENTAL SEED CROPS (FLOWERS, BULBS, ETC.)** **[EXCEPT IN THE STATE OF CALIFORNIA]**

For pre-harvest desiccation of ornamental seed crops. NOT FOR FOOD OR FIBER CROPS.

**Broadcast (Air or Ground):** Apply 1.5-2 pints of this product plus the labeled rate of a 75% or greater nonionic surfactant per acre in sufficient water (minimum of 5 gallons by air; 15 gallons by ground) for desiccation and weed burndown. Repeat as needed at no less than at 5-day intervals; up to three applications. Do not use seed, screenings, or waste as feed or for consumption.

### **USE IN LANDSCAPE, INDUSTRIAL, RECREATIONAL, COMMERCIAL, RESIDENTIAL, AND PUBLIC AREAS**

This product provides fast control of broadleaf and grassy weeds in industrial, recreational, golf course, commercial, residential, and public areas. This product is a nonselective herbicide that rapidly kills undesirable above ground weed growth in 24-36 hours. Avoid applications to desirable plants.

This product is a contact/desiccant herbicide; it is essential to obtain complete coverage of the target weeds to get good control. Improper application technique and/or application to stressed weeds may result in unacceptable weed control. For best results, apply to actively growing,

young weeds. Difficult weeds (such as perennial or deeply-rooted weeds) can often be controlled by tank mixing this product with other systemic-type herbicides. Refer to other product labels for specific application directions.

For residual weed control, tank mix this product with a pre-emergent herbicide labeled for the intended use site. When mixing this product with another herbicide, it is recommended to mix just a small amount to first determine if the mixture is physically compatible before proceeding with larger volumes.

SePRO Corporation has not tested all possible tank mixtures with other herbicides for compatibility, efficacy or other adverse effects. Before mixing with other herbicides SePRO Corporation recommends you first consult your state experimental station, state university or extension agent.

- **Grounds maintenance weed control:** This product can be used as a spot or broadcast spray to control weeds in public, commercial and residential landscapes, including landscape beds, lawns, golf courses and roadsides. This product can also be used for weed control around the edges and non-flooded portions of ponds, lakes and ditches.
- **Trim and Edge weed control:** This product can be used to eliminate undesired grass and broadleaf plant growth in a narrow band along driveways, walkways, patios, cart paths, fence lines, and around trees, ornamental gardens, buildings, other structures, and beneath noncommercial greenhouse benches. Vegetation control with this product is limited to the spray application width. Do not exceed the labeled rate of this product as excessive rates may result in staining of concrete-based materials.
- Since this product does not translocate systemically, can be used as an edging or pruning tool when precisely applied to select areas of grass or to undesirable growth on desirable ornamental bedding plants, ground covers, etc.
- **Industrial weed control:** This product can be used as a spot or broadcast spray either alone or in combination with other herbicides as a fast burndown or control weeds in rights-of-ways, railroad beds/yards, highways, roads, dividers and medians, parking lots, pipelines, pumping stations, public utility lines, transformer stations and substations, electric utilities, storage yards, and other non-crop areas.

**Spot spray:** Apply either 1-2 quarts of this product plus the labeled rate of a 75% or greater nonionic surfactant per 100 gallons water, or 0.75 ounces (22 milliliters) this product plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon of water.

**Broadcast:** 1-2 pints of this product per acre in sufficient water to insure good spray coverage. Add the labeled rate of 75% or greater nonionic surfactant per 100 gallons spray mixture. Greater water volumes are necessary if the target plants are tall and/or dense. It is recommended that 60 gallons or greater water volume be used to obtain good coverage of dense weeds.

#### **USE IN TURF RENOVATION (ALL TURF AREAS EXCEPT COMMERCIAL SOD FARMS)**

To desiccate golf course turf and other turf areas prior to renovation, apply 1-2 pints of this product per acre plus the labeled rate of a 75% or greater nonionic surfactant in 20-100 gallons of water (4 teaspoons of this product plus the labeled rate of a 75% or greater nonionic surfactant per 1 gallon of water) using ground spray equipment. Apply for full coverage and thorough

contact with the turfgrass. Apply only when the turf is dry, free from dew and incidental moisture. For enhanced turf desiccation, especially in the case of thick turfgrass, water volumes should approach 100 gallons of water per acre.

For suppression of regrowth and quick desiccation of treated turfgrass, this product may be mixed with other systemic nonselective or systemic post-emergence grassy weed herbicides. Refer to other product labels for specific application directions and restrictions.

Avoid spray contact with, or spray drift to, foliage of ornamental plants or food crops. Do not graze livestock on treated turf or feed treated thatch to livestock.

### **USE IN DORMANT ESTABLISHED TURFGRASS (BERMUDAGRASS, ZOYSIAGRASS), NONFOOD OR FEED CROP**

For control of emerged annual broadleaf and grass weeds, including little barley<sup>†</sup>, annual bluegrass, bromes including rescuegrass, six-weeks fescue, henbit, buttercup, and Carolina geranium in established dormant Bermudagrass lawns, parks, golf courses, etc.

Apply 1-2 pints this product per acre in 20-100 gallons of spray mix by ground as a broadcast application. Add the labeled rate of a 75% or greater nonionic surfactant per 100 gallons of spray mixture.

Bermudagrass must be dormant at application. Application to actively growing Bermudagrass or Bermudagrass in transition may cause delay or permanent injury. Users in the extreme Southern areas should be attentive to the extent of dormancy at the time of application.

<sup>†</sup>For control of little barley, apply this product prior to the mid-boot stage.

### **USE IN AQUATIC AREAS**

**New York - Not for sale or use in New York State without Supplemental Special Local Needs Labeling.**

**Obtain Required Permits:** Consult with appropriate state or local pesticide and/or water authorities before applying this product in or around public waters. Permits and posting or treatment notification may be required by state, tribal, or local public agencies.

Treatment of dense weed areas may result in oxygen loss from decomposition of dead weeds. This loss of oxygen may cause fish suffocation. Therefore, to minimize this hazard, do not treat more than  $\frac{1}{2}$  of the water body area at one time and wait 14 days between treatments when susceptible plants are mature and have grown to the water's surface, or when the treatment would result in significant reductions in total plant biomass. Waters having limited and less dense weed infestations may not require partial treatments.

For application only to **still water** (i.e. ponds, lakes, and drainage ditches) where there is minimal or no outflow to public waters.

and/or

For applications to **public waters** in ponds, lakes, reservoirs, marshes, bayous, drainage ditches, canals, streams, rivers, and other slow-moving or quiescent bodies of water for control of aquatic weeds. For use by:

- Corps of Engineers;



- Federal or State public agencies (i.e., Water Management District personnel, municipal officials); or
- Applicators and/or licensees (certified for aquatic pest control) that are authorized by the State or Local government.

Treated water may be used according to the water use restrictions set forth in Table 1 or when an approved assay or analytical method establishes that the water does not contain more than the designated maximum contaminant level goal (MCLG) of 0.02 mg/l (ppm) of diquat dibromide (calculated as the cation).

<b>Application Rate (gallons/surface acre)</b>	<b>Drinking</b>	<b>Fishing and Swimming</b>	<b>Livestock/ Domestic Animals Consumption</b>	<b>Irrigation to Turf and Landscape Ornamentals<sup>††</sup></b>	<b>Irrigation to Food Crops and Production Ornamentals<sup>††</sup></b>
≥ 2	3 days	0	1 day	3 days	5 days
1	2 days	0	1 day	2 days	5 days
0.75	2 days	0	1 day	2 days	5 days
0.50	1 day	0	1 day	1 day	5 days
Spot Spray <sup>†</sup> (< 0.5)	1 day	0	1 day	1 day	5 days

<sup>†</sup> Add a nonionic surfactant (with at least 75% of the constituents active as a spray adjuvant) at the rate recommended by the manufacturer.

<sup>††</sup> For preparing agricultural sprays for food crops, turf or ornamentals (to prevent phytotoxicity), do not use water treated with this product before the specified time period.

When the contents of more than one spray tank is necessary to complete a single aquatic application, no water holding restrictions apply between the consecutive spray tanks.

No applications are to be made in areas where commercial processing of fish, resulting in the production of fish protein concentrate or fish meal, is practiced.

**Floating and Marginal Weed Control**

This product may be applied by backpack, airboat, spray handgun, helicopter, airplane, or similar application equipment that results in thorough spray coverage.

cattails, *Typha* spp.  
 duckweed, including *Lemna* spp.  
 frog’s bit<sup>†</sup>, *Limnobium spongia*  
 pennywort, *Hydrocotyle* spp.

salvinia spp., including *Salvinia molesta*  
 water hyacinth, *Eichhornia crassipes*  
 water lettuce, *Pistia stratiotes*

<sup>†</sup>Not for use in California

**Spot Treatment:** Apply this product at 2 to 4 quarts per 100 gallons spray carrier (0.5 - 1.0% solution) with an approved aquatic surfactant or wetting agent at 0.25 - 1.0% v/v (1 quart to 1 gallon per 100 gallons water; refer to the surfactant label for product-specific rates). For cattail control, this product should be applied prior to flowering at the maximum application rate (8 quarts of this product /100 gallons spray carrier) plus the wetting agent. Repeat treatments may be necessary for complete control.

Spray to completely wet target weeds but not to runoff. Densely packed weeds or mats may require additional applications due to incomplete spray coverage. Re-treat as needed. For best results, re-treat weed escapes within 2 weeks of the initial treatment.

**Broadcast Treatment:** Apply this product at the rate of 0.5 - 2.0 gallons per surface acre in sufficient carrier along with 16-32 ounces per acre of an aquatic surfactant or wetting agent (refer to the surfactant label for product specific rates). Re-treat as necessary for densely populated weed areas. Good coverage is necessary for control of the target weeds.

For duckweed control, apply this product at 1 - 2 gallons/acre.

### Submersed Weed Control

To control submersed weeds apply this product in water at 0.5 - 2.0 gallons per surface acre (per 4 foot water depth), or up to 0.5 gallons/acre foot in water with an average depth greater than 4 feet deep. For severe weed infestations or when treating more difficult to control species, use 0.5 gallons/acre foot of water. Refer to Table 2 for application rates.

algae <sup>††</sup> , <i>Spirogyra</i> spp. and <i>Pithophora</i> spp.	hydrilla, <i>Hydrilla verticillata</i>
bladderwort, <i>Utricularia</i> spp.	naiads, <i>Najas</i> spp.
Brazilian elodea, <i>Egeria densa</i>	pondweeds <sup>†</sup> , <i>Potamogeton</i> spp.
coontail, <i>Ceratophyllum demersum</i>	watermilfoils, including Eurasian,
elodea, <i>Elodea</i> spp.	<i>Myriophyllum</i> spp.

<sup>†</sup>This product controls *Potamogeton* species except Richardson's pondweed, (*P. richardsonii*).  
<sup>††</sup>Suppression only. For control of *Spirogyra* and/or *Pithophora*, use this product in a tank mix with an approved algaecide.

TABLE 2. GALLONS OF PRODUCT PER SURFACE ACRE <sup>†</sup>				
Application Rate (gallons/acre)	Average Water Depth			
	1 Foot	2 Feet	3 Feet	4 Feet <sup>††</sup>
1	0.25 gal.	0.50 gal	0.75 gal.	1.0 gal.
2	0.50 gal	1.0 gal.	1.5 gals.	2.0 gals.

<sup>†</sup>For water depths ≤ 2 feet including shorelines, do not exceed 1 gallon per surface acre.  
<sup>††</sup>In treatment areas with an average water depth greater than 4 feet, apply a maximum of 0.5 gallons per acre foot of water.

**Subsurface Applications:** Where the submersed weed growth, especially hydrilla, has reached the water surface, apply either in a water carrier or an invert emulsion through trailing hoses to apply the dilute spray below the water surface to insure adequate coverage.

**Bottom Placement:** Where submersed weeds such as hydrilla, bladderwort, or coontail are growing in deeper water and are less mature (e.g. not to the surface of the water) and/or where the water is slowly moving through the weed growth, the use of an application method (such as invert emulsion carrier or long-trailing hoses) to inject this product near the bottom with weighted hoses may improve control.

**Surface Application for Submerged Aquatic Weeds:** Apply the recommended rate of this product as a spray in sufficient carrier to fully cover the target area. Applications should be made to ensure complete coverage of the weed areas. In mixed weed populations, use the high rate of application as indicated by weeds present. For dense submersed weeds or water over 2 feet

deep, a surface spray is not recommended (This product should be applied subsurface in these situations.)

#### **TANK MIXES WITH OTHER AQUATIC HERBICIDES/ALGAECIDES:**

For severe weed or algae infestations, the use of an approved algaecide either as a pretreatment to the application of this product or in a tank mix, may result in enhanced weed control.

When tank mixing, read and follow the labeled precautionary statements, directions for use, weeds controlled, and other restrictions for each tank mix product. Use in accordance with the most restrictive label limitations and precautions of the products used in the tank mix. Do not exceed any labeled rate or dose. To ensure compatibility, a jar test is recommended before field application of any tank mix combination. Consult with SePRO Corporation for latest tank mix recommendations.

#### **Harvester + Komeen<sup>®</sup>**

The addition of Komeen, or other copper-based herbicides/algaecides, with this product may improve control on some species, such as hydrilla. For best results, apply 2 gallons this product in combination with 4 gallons of Komeen (0.8 lbs. a.i./gallon) per acre. For hydrilla control and control of other species with high sensitivity to copper, lower rates of Komeen may also enhance the activity of this product. Apply copper at a minimum of 0.1 ppm in combination with this product. Higher rates may be needed in areas with dense weeds.

#### **Harvester + endothall**

The addition of endothall with this product may improve control on some species, such as hydrilla. For best results, apply this product at 1 to 2 gallons per acre in combination with the dipotassium salt of endothall at 0.6 to 1.2 gallons/acre foot (i.e. 1 to 2.0 ppm a.i.). Higher rates may be used, but do not exceed the maximum allowed rate for either product.

#### **NOTE: For Drinking (Potable) Water**

The drinking (potable) water restrictions for applications of this product plus endothall are to ensure that consumption of water by the public is allowed only when the concentration of endothall in the water is less than the MCL (Maximum Contamination Level) of 0.1 ppm. Applicators should consider the unique characteristics of the treated waters to assure that endothall concentrations in potable drinking water do not exceed 0.1 ppm at the time of consumption.

For applications of this product plus endothall, the drinking water setback distance from functioning potable water intakes is  $\geq 600$  feet. Note: Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Do not contaminate feed, foodstuffs, or drinking water. Do not store or transport near feed or food. Store at temperatures above 32°F.

**Pesticide Disposal:** Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Non-refillable Container Disposal (rigid, 5 gallons or less):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer the container for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Non-refillable Container Disposal (rigid, greater than 5 gallons):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling (if available) or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.


**Container Handling (bulk): Triple rinse (or equivalent).** Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

**Warranty Disclaimer:** SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

**Misuse:** Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <http://seprolabels.com/terms> or scan the image below.



© Copyright 2022 SePRO Corporation

® Harvester is a registered trademark of SePRO Corporation

EPA Accepted Date 11/15/2019

FPL20191115

SePRO Corporation 11550 N. Meridian St., Suite 600 Carmel, IN 46032 U.S.A.