

Camelot* 0 Fungicide/Bactericide



FOR ORGANIC PRODUCTION



Special Solutions. For Specialty Markets

Use Sites:

Greenhouses **Shadehouses Outdoor nurseries** Outdoor landscape plantings Interior landscapes **Turfgrass**

Formulation:

10% Flowable Liquid

Packaging:

1 gallon container

Active Ingredient:

Copper Octanoate (Copper Soap)

Chemical Class:

Copper Complex

Mode of Action (FRAC Code):

Multi-site (M1)

Rate:

0.5 to 2.0 gallons per 100 gallons (64 to 256 fl. oz. per 100 gallons)

Restricted Entry Interval (REI):

4 hours

Application Technique:

Apply as a thorough spray on all plant surfaces. Spray foliage and stems to run-off.

Application Timing:

Initiate applications at first sign of disease or when conditions are favorable for disease development. If necessary, repeat applications should be made at 7 to 10 day intervals.

The Product for Complete Disease Control – Now for **Organic Production**

Camelot O is a liquid copper formulation using proven technology to provide broad-spectrum bacterial and fungal disease control on a wide variety of ornamental plants, greenhouse vegetable/herb crops and turfgrass. Camelot O is a uniquely patented formula of fixed copper combined with fatty acids to form a true soap providing performance advantages over other copper-based products.



Camelot O requires minimal elemental copper for excellent disease control through its superior coverage ability and foliar adherence. Camelot O gives you the complete disease control you want and deserve.

Features of Camelot O Fungicide/Bactericide are:

- Broad-spectrum fungal and bacterial disease control
- Multiple modes of action for resistance management
- · Control of bacterial leaf spots and blights
- Approved for Organic Production by the National Organic Program (NOP)
- Easy to measure and use flowable copper soap formulation
- Use on greenhouse vegetable and herb crops
- · 4-hour REI

Diseases Controlled

Camelot O controls a broad-spectrum of ornamental fungal and bacterial pathogens.

Bacterial Pathogens

Erwinia spp.

Pseudomonas spp.

Xanthomonas spp.

Bacterial Leaf Spot and Blight

Fungal Pathogens

- Anthracnose
- Ascochyta Leaf Blight
- Black Spot
- Botrytis Blight (Gray Mold)
- Cercospora Leaf Blight/Spot
- Dollar Spot
- · Downy Mildew
- Leaf Spots
- Needle Blights and Casts
- Powdery Mildew
- Rusts
- · White Mold



Listed by the Organic Materials Review Institute (OMRI) for use in organic production.


1-800-419-7779 www.sepro.com

Application Information


Use Camelot O in a disease control program at 0.5 to 2.0 gallons per 100 gallons (64 to 256 fl. oz./100 gallons) and apply to one (1) acre. Initiate applications at first sign of disease or when conditions are favorable for disease development. If necessary, repeat applications should be made at 7 to 10 day intervals. A spreader may be used to improve the spreading of Camelot O on hard to wet plants.

Tank-mix Restrictions

Do not tank-mix with chelated or liquid fertilizers.



Dr. A.R. Chase, Chase Research Gardens, Inc., 2010.



Dr. A.R. Chase, Chase Research Gardens, Inc., 2010.



Fungicide/Bactericide